

## Nuclear Spin Dependence of Hydrogenic Plasmas in the Laboratory and the Diffuse Interstellar Medium

Kyle N. Crabtree, *Department of Chemistry, University of Illinois, Urbana, IL 61801*; Nick Indriolo, *Department of Astronomy, University of Illinois, Urbana, IL 61801*; Holger Kreckel, Carrie A. Kauffman, Brian A. Tom, Eftalda Bećka, Brett A. McGuire, *Department of Chemistry, University of Illinois, Urbana, IL 61801*; Benjamin J. McCall, *Departments of Chemistry, Astronomy, and Physics, University of Illinois, Urbana, IL, 61801*.

Observations of diffuse molecular clouds have shown that the excitation temperature  $T(\text{H}_3^+)$  derived from the  $(J, K) = (1, 0)$  (*ortho*) and  $(1, 1)$  (*para*) rotational levels of  $\text{H}_3^+$  does not necessarily agree with the kinetic temperature ( $T_{01}$ ) inferred from UV measurements of  $\text{H}_2$ . In four of the five diffuse molecular cloud sight lines for which both  $\text{H}_3^+$  and  $\text{H}_2$  observations are available,  $T(\text{H}_3^+)$  is lower than  $T_{01}$  by 30 K. The reaction  $\text{H}_3^+ + \text{H}_2 \rightarrow \text{H}_2 + \text{H}_3^+$  is expected to thermalize the  $\text{H}_3^+$  nuclear spin distribution, but the interplay of nuclear spin selection rules and energetic restrictions at the low temperatures of the diffuse interstellar medium may prevent full thermalization at steady state. Alternatively, the nonthermal distribution could arise if  $\text{H}_3^+$  does not experience a sufficient number of thermalizing collisions with  $\text{H}_2$  during its lifetime.

We have studied the nuclear spin dependence of the reaction of  $\text{H}_3^+$  with  $\text{H}_2$  in the laboratory by measuring the *ortho:para* ratio of  $\text{H}_3^+$  formed in plasmas of varying *ortho:para*  $\text{H}_2$  ratios. This study was performed in a hollow cathode cell which enabled the first measurements of this reaction at low temperature (130 K). From these measurements, we derived the ratio of the “proton hop” and “hydrogen exchange” rate coefficients as a function of temperature. The ratio,  $\alpha$ , was found to decrease with temperature, already reaching the statistical limit of 0.5 at a temperature of 130 K.

Knowledge of  $\alpha$  enables modeling of the *ortho:para* ratio of  $\text{H}_3^+$  in diffuse molecular clouds. We have modeled the nuclear spin dependence of the formation, thermalization, and destruction processes of  $\text{H}_3^+$ , and found that the nonthermal distribution is not caused by nuclear spin selection rules or energetic restrictions in the  $\text{H}_3^+ + \text{H}_2$  reaction. Rather, it is likely caused by incomplete thermalization prior to  $\text{H}_3^+$  destruction via electron dissociative recombination.